

NCCN Guidelines Version 2.2025 Myeloproliferative Neoplasms

NCCN Guidelines Index
Table of Contents
Discussion

MYELOPROLIFERATIVE NEOPLASM SYMPTOM ASSESSMENT FORM: TOTAL SYMPTOM SCORE (MPN-SAF TSS)¹

(Recommended for monitoring symptoms during the course of treatment)

Symptom	1 to 10 (0 if absent) ranking 1 is most favorable and 10 least favorable
Please rate your fatigue (weariness, tiredness) by circling the one number that best describes your WORST level of fatigue during past 24 hours	(No Fatigue) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)

Circle the one number that describes, during the past week, how much difficulty you have had with each of the following symptoms

Filling up quickly when you eat (early satiety)	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Abdominal discomfort	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Inactivity	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Problems with concentration- compared to prior to my MPD	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Night sweats	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Itching (pruritus)	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Bone pain (diffuse not joint pain or arthritis)	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Fever (>100 F)	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Daily)
Unintentional weight loss last 6 months	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)

Note: All recommendations are category 2A unless otherwise indicated.

¹ Reproduced with permission from Emanuel RM, Dueck AC, Geyer HL, et al. Myeloproliferative neoplasm (MPN) symptom assessment form total symptom score: prospective international assessment of an abbreviated symptom burden scoring system among patients with MPNs. J Clin Oncol 2012;30:4098-4103.